

## VU Research Portal

### **“Movement of the sacroiliac joint during the active straight leg raise test in patients with long-lasting severe sacroiliac joint pain”**

Hu, Hai; Prins, Maarten R.; Wu, Wen Hua; van Dieën, Jaap H.; Xia, Chun; Mens, Jan M.A.; Meijer, Onno G.

#### ***published in***

Clinical Biomechanics  
2018

#### ***DOI (link to publisher)***

[10.1016/j.clinbiomech.2018.01.008](https://doi.org/10.1016/j.clinbiomech.2018.01.008)

#### ***document version***

Publisher's PDF, also known as Version of record

#### ***document license***

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

#### ***citation for published version (APA)***

Hu, H., Prins, M. R., Wu, W. H., van Dieën, J. H., Xia, C., Mens, J. M. A., & Meijer, O. G. (2018). “Movement of the sacroiliac joint during the active straight leg raise test in patients with long-lasting severe sacroiliac joint pain”: A letter to the editor. *Clinical Biomechanics*, 52, 100-101.  
<https://doi.org/10.1016/j.clinbiomech.2018.01.008>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)



## Letter to the Editor

**“Movement of the sacroiliac joint during the active straight leg raise test in patients with long-lasting severe sacroiliac joint pain”—A letter to the editor**



## ARTICLE INFO

## Keywords:

Active straight leg raise  
Pelvic girdle pain  
Sacroiliac joints  
Pubic symphysis

Dear Editor of *Clinical Biomechanics*,

We read with interest Kibsgård et al.'s above *Clinical Biomechanics* paper. Radiostereometric Analysis was performed to measure sacroiliac movements during the Active Straight Leg Raise (ASLR) in 12 patients with chronic Pelvic Girdle Pain (PGP). Of the patients, 8 had been diagnosed with bilateral, and 4 with unilateral sacroiliac pain. All were about to undergo sacroiliac fusion surgery.

We fully agree with Kibsgård et al. that this is an important field. About 5% of all pregnant women suffer from PGP that is sufficiently serious to warrant medical attention (Wu et al., 2004). Also athletes with groin pain may have PGP (Verrall et al., 2001), often in the form of osteitis pubis. Both groups of patients have difficulties performing the ASLR (Mens et al., 2006).

Contralateral to the side of the raising leg, Kibsgård et al. observed  $0.8^\circ$  backward rotation of the ilium with respect to the sacrum (range  $-0.3^\circ$  to  $-1.3^\circ$ , ‘minus’ for backward), and ipsilaterally on average  $0.0^\circ$  (range  $-1.0^\circ$  to  $+0.5^\circ$ ). We congratulate Kibsgård et al. with this contribution to the field. Still, there are some questions.

In attempting to understand the causes of the observed movements, Kibsgård et al. invoke our earlier ASLR study, Hu et al. (2012), with healthy subjects, where we reasoned that hip flexion exerts a forward pull on the ipsilateral ilium, which may be counteracted by contralateral extensor activity. This may only work “if the two sides of the pelvis act as a single unit, such as when they are pressed together by force closure” (Hu et al., 2012, p. 534; cf. Snijders et al., 1983). But Kibsgård et al.'s contralateral sacroiliac movement revealed that force closure, if any, was not, or not completely successful. Perhaps, failing force closure differentiates PGP from healthy subjects, but presently, we cannot know this, since Hu et al. (2012), did not measure sacroiliac movements, and Kibsgård et al. (2017), did not use healthy controls.

And how is the ipsilateral ilium kept together with the sacrum? Kibsgård et al. invoke ipsilateral-only stabilizing mechanisms, but until now, such mechanisms remain unknown. Kibsgård et al. remark that the ilium may already be in maximum anterior rotation when the subject is in supine position (Sturesson et al., 1989), which would not allow any further forward rotation. Still, to understand what happens, we need to know the initial position, and the movement, if any, of both ilia and the sacrum, and not just the relative movement of the ilium with respect to the sacrum.

Mens et al. (1999), reported 1–5 mm vertical movements (“steps”) between the pubic bones, with the ipsilateral pubis lower than the contralateral one. They explained this as a forward rotation of the ipsilateral ilium (cf. Sturesson et al., 1989). Kibsgård et al. state that they “contradict” (p. 44) this interpretation, which difference of opinion, they argue, may be “due to different methodology”, their own measurements being “more reliable and valid” (p. 43). But we see no role of methodology here.

Mens et al. did not measure sacroiliac movements, but inferred them from symphyseal steps, whereas Kibsgård et al. did not measure symphyseal steps, but found sacroiliac movements that may affect these steps. Assuming rigidity of the pelvic bones (but see Haussler et al., 2009), we performed a quick and dirty calculation, which revealed that a  $0.8^\circ$  backward rotation of the contralateral ilium implies in the order of 2 mm stepping upward of the pubis, which, ballpark, agrees with Mens et al. Any asymmetry of sacroiliac sagittal plane movements will lead to symphyseal steps.

Comparing painful and pain free joints, Kibsgård et al. found significantly larger rotations around the helical axes of the painful joints,  $0.8^\circ$ , than in the pain free joints,  $0.4^\circ$ . Thus, more movement may coincide with more pain. In terms of cartesian rotation with respect to the sacrum, however, the authors did not find any systematic differences, as if there is no relationship between sacroiliac movement and pain. Are we missing something important here, such as, perhaps, pain upon sacroiliac compression (Szadek et al., 2010)? We wonder if there was any relationship at all between individual ASLR scores and sacroiliac movements (helical/cartesian; ipsilateral/contralateral). Was there more pain in subjects with more forward rotation of the ipsilateral ilium?

Further understanding of ASLR problems in PGP remains an important priority. The next step in research should include full analyses of muscle activity, pelvic kinematics, the patients' problems, and the relationships between these variables.

\* Support from the National Natural Science Foundation is acknowledged by H. Hu (#31270996) and W.-H. Wu (#81272161).

## We need to better understand the Active Straight Leg Raise

## References

- Haussler, K.K., McGilvray, K.C., Ayturk, U.M., Puttlitz, C.M., Hills, A.E., McIlwraith, C.W., 2009. Deformation of the equine pelvis in response to in vitro 3D sacroiliac joint loading. *Equine Vet. J.* 41, 207–212.
- Hu, H., Meijer, O.G., Hodges, P.W., Bruijn, S.M., Strijers, R.L., Nanayakkara, P.W., Van Royen, B.J., Wu, W.H., Xia, C., Van Dieën, J.H., 2012. Understanding the Active Straight Leg Raise (ASLR): an electromyographic study in healthy. *Man. Ther.* 17, 531–537.
- Kibsgård, T.J., Röhrli, S.M., Røise, O., Stuessen, B., Stuge, B., 2017. Movements of the sacroiliac joint during the Active Straight Leg Raise test in patients with long-lasting severe sacroiliac joint pain. *Clin. Biomech.* 47, 40–45.
- Mens, J.M., Vleeming, A., Snijders, C.J., Stam, H.J., Ginai, A.Z., 1999. The active straight leg raising test and mobility of the pelvic joints. *Eur. Spine J.* 8, 468–473.
- Mens, J., Inklaar, H., Koes, B.W., Stam, H.J., 2006. A new view on adduction-related groin pain. *Clin. J. Sports Med.* 16, 15–19.
- Snijders, C.J., Vleeming, A., Stoeckart, R., 1983. Transfer of lumbosacral load to iliac bones and legs, part 1: biomechanics of self-bracing of the sacroiliac joints and its significance for treatment and exercise. *Clin. Biomech.* 8, 285–294.
- Stuessen, B., Selvik, G., Udén, A., 1989. Movements of the sacroiliac joints: a roentgen stereophotogrammetric analysis. *Spine* 14, 162–165.
- Szadek, K.M., Hoogland, P.V., Zuurmond, W.W., De Lange, J.J., Perez, R.S., 2010. Possible nociceptive structures in the sacroiliac joint cartilage: an immunohistochemical study. *Clin. Anat.* 23, 192–198.
- Verrall, G.M., Slavotinek, J.P., Fon, G.T., 2001. Incidence of pubic bone marrow oedema in Australian rules football players: relation to groin pain. *Br. J. Sports Med.* 35, 28–33.
- Wu, W.H., Meijer, O.G., Mens, Uegaki K., JM, Van Dieën J.H., Wuisman, P.I., Östgaard, H.C., 2004. Pregnancy-related pelvic girdle pain (PPP), I: terminology, clinical presentation, and prevalence. *Eur. Spine J.* 13, 575–589.

Hai Hu<sup>a</sup>, Maarten R. Prins<sup>b</sup>, Wen-Hua Wu<sup>c,\*</sup>, Jaap H. van Dieën<sup>d</sup>, Chun Xia<sup>e</sup>, Jan M.A. Mens<sup>f</sup>, Onno G. Meijer<sup>d,g</sup>

<sup>a</sup> Department of Orthopaedic Surgery, Affiliated Sixth People's Hospital, Jiao Tong University, Shanghai, PR China

<sup>b</sup> Military Rehabilitation Centre “Aardenburg”, Doorn, The Netherlands

<sup>c</sup> Department of Orthopaedic Surgery, Second Affiliated Hospital, Fujian Medical University, Zhongshan North Road 34, Quanzhou 362000, Fujian, PR China

<sup>d</sup> Department of Human Movement Sciences, Vrije Universiteit, Amsterdam, The Netherlands

<sup>e</sup> Department of Joint Surgery & Sports Medicine, Zhongshan Hospital, Xiamen University, Xiamen, Fujian, PR China

<sup>f</sup> Department of Rehabilitation Medicine & Physical Therapy, Erasmus University Medical Center, Rotterdam, The Netherlands

<sup>g</sup> Orthopaedic Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China

E-mail address: drwuwenhua@126.com

\* Corresponding author at: Orthopaedic Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China.